





















































































LATIN NAME	COMMON NAME	ZONE	SOIL	EXPOSITION	BLOOMING							SOWING METHOD	
					APRIL	MAY	JUNE	JULY	AUG.	SEPT.	OCT.		
Acer saccharum	Sugar Maple	3	cool										2
Juglans nigra	Black walnut	3	cool										11
Quercus rubra	Red Oak	3	cool or moist										11
Aronia melanocarpa	Black Chokeberry	2	cool or moist or water soaked										2
Cephalanthus occidentalis	Button-bush	4	moist or water soaked										2
Cornus alternifolia	Alternate-leaved Dogwood	3	cool or moist										9
Cornus stolonifera	Red-osier Dogwood	1	cool or dry										9
Ilex verticillata	Winterberry	3	cool or moist or water soaked										2
Physocarpus opulifolius	Nine-bark	4	cool or dry										6
Prunus virginiana	Choke Cherry	2	cool or dry										7
Rhus typhina	Vinegar-tree	3	cool or dry										12
Sambucus canadensis	Canadian Elder	3	cool										9
Viburnum trilobum	Cranberry-tree	2	cool										9
Calamagrostis canadensis	Canada Reed-grass	3	cool or moist										2
Hierochloe odorata	Vanilla Grass	3	cool										4
Celastrus scandens	Climbing Bittersweet	4	cool										10
Clematis virginiana	Virginia Virgin's Bower	3	cool										6
Echinocystis lobata	Wild Cucumber	3	cool or moist										10
Parthenocissus quinquefolia	Virginia Creeper	2	cool										6
Vitis riparia	Wild Grape	2	Cool ,moist or dry										6
Anaphalis margaritacea	Life-everlasting	3	dry										8
Aquilegia canadensis	Wild Columbine	3	cool and dry										8
Asclepias incarnata	Swamp Milkweed	3	cool or moist or water soaked										5
Aster novae-angliae	New England Aster	3	cool and dry										1
Epilobium angustifolium	Firewood	2	cool and dry										3
Heliopsis helianthoides	False Sunflower	4	cool										2
Hypericum pyramidatum	Giant St.John's-wort	4	cool and moist										8
Iris versicolor	Larger Blue-Flag	2	cool or moist or water soaked										12
Lilium canadense	Wild Yellow Lily	4	cool or moist										1
Monarda fistulosa	Wild Bergamot	4	cool										3
Oenothera victorinii	Victorin's Evening Primrose	2	cool or dry										12
Verbena hastata	Blue Vervein	4	cool or moist or water soaked										8

- 1) No pre-treatment necessary. Sow seeds when there is no risk of ground frost.
- 2) No pre-treatment necessary. Sow seeds when there is no risk of ground frost. Or plant the seeds outdoors during the fall. Germination will occur the following spring or later on.
- 3) No pre-treatment necessary. Sow seeds when there is no risk of ground frost. Sow seeds on the surface without covering after sowing. These seeds are very small and need light to germinate. Seeds can be mixed with fine sand or talcum for a uniform sowing.
- 4) Mix seeds with an equal amount of damp sand or vermiculite. Place the mixture in a sealed plastic bag. Store the bag in a refrigerator or cool storage room for 30 days. Sow seeds when there is no risk of ground frost.
- 5) Mix seeds with an equal amount of damp sand or vermiculite. Place the mixture in a sealed plastic bag. Store the bag in a refrigerator or cool storage room for 30 days. Sow seeds when there is no risk of ground frost. Or plant the seeds outdoors during the fall. Germination will occur the following spring or later on.
- 6) Mix seeds with an equal amount of damp sand or vermiculite. Place the mixture in a sealed plastic bag. Store the bag in a refrigerator or cool storage room for 30 to 90 days. Sow seeds wherever desired as soon as the soil gets warm enough. Or plant the seeds outdoors during the fall. Germination will occur the following spring or later on.
- 7) Mix seeds with an equal amount of damp sand or vermiculite. Place the mixture in a sealed plastic bag. Store the bag in a refrigerator or cool storage room for 90 days. Sow seeds when there is no risk of ground frost. Or plant the seeds outdoors during the fall. Germination will occur the following spring or later on.
- 8) Mix seeds with an equal amount of damp sand or vermiculite. Place the mixture in a sealed plastic bag. Store the bag in a refrigerator or cool storage room for 30 to 90 days. Sow seeds when there is no risk of ground frost. Sow seeds on the surface without covering after sowing. These seeds are very small and need light to germinate. Seeds can be mixed with fine sand or talcum for a uniform sowing.
- 9) Mix seeds with an equal amount of damp sand or vermiculite. Place the mixture in a sealed plastic bag. Store the bag in a refrigerator or cool storage room for 90 days. Sow seeds when there is no risk of ground frost. Or plant the seeds outdoors during the fall. Germination will occur after 2 winter periods (18 to 20 months).
- 10) Bring water to a boil for 5 minutes. Pour over seeds and let soak for 24 hours. Mix seeds with an equal amount of damp sand or vermiculite. Place the mixture in a sealed plastic bag. Store the bag in a refrigerator or cool storage room for 90 days. Sow seeds when there is no risk of ground frost. Or plant the seeds outdoors during the fall. Germination will occur the following spring or later on.
- 11) Do not allow the seeds to dry. Bring water to a boil for 5 minutes. Pour over seeds and let soak for 24-48 hours. Mix seeds with an equal amount of damp sand or vermiculite. Place the mixture in a sealed plastic bag. Store the bag in a refrigerator or cool storage room for 30 to 90 days. Sow seeds when there is no risk of ground frost. Or plant the seeds outdoors during the fall. Germination will occur the following spring or later on.
- 12) Lightly abrade the seed coats by rubbing seeds between two sheets of sandpaper. Bring water to a boil for 5 minutes. Pour over seeds and let soak for 24 hours. Mix seeds with an equal amount of damp sand or vermiculite. Place the mixture in a sealed plastic bag. Store the bag in a refrigerator or cool storage room for 90 days. Sow seeds when there is no risk of ground frost. Or, plant the seeds outdoors during the fall. Germination will occur the following spring or later on.